Your Guide to

Colonoscopy Suprep





Medications that you will need

A prescription for Suprep will be sent to your pharmacy. Please follow the directions below when taking your Suprep. If your pharmacy does not receive this prescription, please call the office at: ______.

How to take medications for colonoscopy

On the day **BEFORE** your colonoscopy

Only eat foods listed below on the day before your colonoscopy

- Water
- Coffee with NO creamer
- Tea with NO creamer
- Broth (beef, chicken or vegetable) (no meat, noodle or vegetable pieces)
- Apple or white grape juice
- Sprite, ginger ale and 7-Up
- Gatorade or PowerAde (no red, blue or purple)
- Popsicles (no red, blue or purple)
- Jell-O (no red, blue or purple)

If you are diabetic, please buy sugar free options

At 6 p.m.

- 1. Prepare your Suprep mixture: Pour one of the brown bottles into the cup provided. Top the cup off with cool water. Save the second bottle for the morning of your colonoscopy.
- 2. Drink the Suprep mixture followed by four (4) cups (32 ounces) of water.

Feelings of nausea and bloating are normal and will subside.

Colonoscopy Suprep, Continued



On the day of your colonoscopy

Five (5) hours before the time of your colonoscopy

Example: If your colonoscopy is scheduled for 9 a.m. you should start drinking your Suprep mixture at 4 a.m. It should be finished by 6 a.m. Do **NOT** drink **ANYTHING** after 7 a.m.

You cannot have anything to eat or drink **EXCEPT** your Suprep mixture on the morning of your colonoscopy.

You can take your medications with a sip of water the day of your colonoscopy unless told not to take them by your doctor.

Be sure to have finished your Clenpiq prep and taken your medications at least two (2) hours before your colonoscopy. You should have **NOTHING** to drink two (2) hours before your colonoscopy.

Procedure Date:	Provider:
Location:	
•	pefore your procedure by Fairfield Medical Center for the londay procedure). Calls will be made sometime prior to

Continue to next page for additional instructions

Colonoscopy Suprep, Continued

Things to remember

- Reminder: You cannot have anything to eat or drink (including water) the day of your procedure except your bowel prep and a small sip of water to take your medicine.
- Someone has to drive you to the hospital, stay at the hospital until you are done and drive you home. You cannot drive the day of your colonoscopy. **If you do not have someone to do these things, your colonoscopy will be cancelled.**
- If you take an iron supplement, stop taking them five (5) days before your procedure to avoid discoloration of your stool.
- The office will let you know which of your medications need to be stopped and for how long. Some of the most commonly adjusted medications include blood thinners and diabetic medications. Medication instructions:

•	
•	
•	

- If drinking the Suprep mixture makes you sick to your stomach, you can try drinking it slower. Some nausea and bloating is normal.
- Your poop should be all liquid. It should not be brown in color. It should look clear to yellow. You will need to reschedule if it does not look like this the morning of your colonoscopy.
- If you need to cancel your colonoscopy you will need to call your doctor's office.

The phone number is: _____



Fairfield Medical Center Endoscopy 740-687-8158 401 N. Ewing St. Lancaster, Ohio 43130 fmchealth.org